

# ISST JOURNAL OF APPLIED CHEMISTRY

Volume 2 No. 1

January – June 2011

## CONTENTS

## Page No.

- Chelating Effect of Multidentate Hybrid Ligands in Heteroleptic Metal Complexes of Heavy Chalcogenides  
*Prarthana Srivastava* 1-4
- Physico-Chemical and Biological Studies on Pollution Potential of River Devaha at District Pilibhit (U.P.) during Winter Season  
*Anoop Chandra, P.N. Saxena, Preeti Saxena and S. Ghazala. Imam* 5-9
- A New Green Catalytic Approach for the Oxidative Cyclization of Nitrogen Sulphur Ligand  
*Sulekh Chandra and Avdhesh Kumar* 11-14
- Comparative Study of the Seasonal Variation of SO<sub>2</sub> Gas in Polluted Air  
*Vandana Srivastava and Abu Shahma* 15-19
- Development of Nitrate Selective Membrane Sensor using Zinc (II) bis(5-phenyl azo salicylaldehyde)-1,2-Benzene Diimine as Electroactive Material  
*Gyanendra Singh, Sanjay Singh, Arpit Singh and Kalash Chandra Yadav* 21-24
- Effect of Soaking Time on Transparent Base Glaze at Moderate Temperature  
*Neetu Sharma, H. Kaur and L.K. Sharma* 25-30
- Anti-Inflammatory Activity of Seeds Extract of *Prosopis Spicigera*  
*Rajiv Agrawal, H.K. Garg, Udit Garg and S.K. Singh* 31-34
- Organic Reactions in Microwave Oven  
*Okram Mukherjee Singh and L. Ronibala Devi* 35-37
- Synthesis and Characterization of Cerium (IV) Phosphoiodate : A New Inorganic Cation Exchanger  
*Subhash Chand, Seema, Brajesh Pal and Rekha* 39-47
- Application of Supercritical Fluid Technology for Natural Products  
*Bina Rani, Upma Singh and Raaz Maheshwari* 49-54
- A Novel System for Cross Linking Fluoroelastomers  
*Khagesh Kumar Singh and Anju Singh* 55-59



**Intellectuals Society for Socio-Techno Welfare**  
Ghaziabad – 201 003 (U.P.) India  
[www.isst.org.in](http://www.isst.org.in)

# ANTI-INFLAMMATORY ACTIVITY OF SEEDS EXTRACT OF PROSOPIS SPICIGERA

<sup>1</sup> Rajiv Agrawal, <sup>3</sup> H.K. Garg, <sup>2</sup> Udit Garg and <sup>1</sup> S.K. Singh

<sup>1</sup> Department of Chemistry, D.B.S. (P.G.) College, Kanpur - 208006, U.P., India

<sup>2</sup> Department of Microbiology, D.B.S. (P.G.) College, Kanpur - 208006, U.P., India

<sup>3</sup> Department of Chemistry, D.V. (P.G.) College, Orai - 285001, U.P., India

E-mail: dr.rajivagrwal@gmail.com

## ABSTRACT

*Prosopis spicigera* Sm. (Leguminosae) is a commonly used plant in Indian traditional medicine. In the current study anti-inflammatory activity of ethanolic extract of *P. spicigera* Seeds was investigated using different animal models. The extract was also subjected to phyto-chemical analysis and their toxic potential. The anti-inflammatory activity was measured using the carrageenan-induced paw edema test. The air dried seeds of *P. spicigera* were soaked in ethanol (1:4, w/v) at the temperature of 60°C for 72 h. The supernatant collected was evaporated under reduced pressure and kept at 50°C until used. The extract was emulsified using 0.1% Tween -80 in normal Saline to prepare the doses of 33, 100 and 300 mg kg<sup>-1</sup>, before performing each experiment. Anti-inflammatory activity in a dose-dependant manner in all assays used to the extract administered intraperitoneally exhibited significant ( $p < 0.05$ ).

**Key words:** *Prosopis spicigera*, Leguminosae, Seeds, ethanol extracts, toxicity, anti-inflammatory activity

## 1. INTRODUCTION

*Prosopis spicigera* Sm. (Leguminosae) is commonly known as Jhand or Shami [4,6]. Yadav R. N. et. al. (1999) use this plant for Cardenolide, and it is found basically throughout the India but abundantly in Bundelkhand region of Uttar Pradesh (INDIA). It is widely used in the traditional medical practice of people living in India, Lanka and China to treatment of headache, skin diseases, ulcers, uterine trouble along with as a safe guard against miscarriage during pregnancy in women [3], laxative and used as tonic [2], used as a hair removal [8], an emulsion of the bark used as a remedy in rheumatism and scorpion sting, current awareness in flavour and fragrance Journal 2007 inter-science widey.com. Earlier some worker have reported the presence of various biologically active constituents in this plant Jewers K.et.al., phytochemistry (1976).[9,18]

In Bundelkhand region this plant is traditionally used to treat headache, stomach ache and as anti-inflammatory drug [13,14,15]. Since its recognition to pain removal as an important chemical entity, various efforts have been made to find a suitable remedial measure, as the drugs used to reduce acid secretion have dominated the pharmacological basis of pain inflammation therapy [7,12,19]. Keeping in view the frequent folklore use of *P. spicigera*, the present study was carried out to determine the anti-inflammatory activity of *P. spicigera* seeds using animal models.

## 2. MATERIALS AND METHOD

*Plant material and preparation of its extracts:-*

*P. spicigera* seeds were collected locally from the region of Bundelkhand (U.P.), sun dried and grinded into powder form. The powdered seeds (300g) at  $25 \pm 3^\circ\text{C}$  was mixed with absolute alcohol (1:4, w/v) placed in water bath at  $60^\circ\text{C}$  for 72h and then filtered by using whatman no.1 to obtain the supernatant. The collected supernatant was then evaporated up to dryness at  $100^\circ\text{C}$ . under reduced pressure and crude dried extract obtain, labeled as EPSS (ethanolic *Prosopis spicigera* supernatant), was kept at  $5^\circ\text{C}$ .

EPSS was emulsified using 0.1% Tween -80 in normal Saline to the doses of 33, 100, 300 mg Kg<sup>-1</sup> body weight for administration into the animals.

### Preliminary phytochemical screening

The alcoholic extract was subjected to preliminary phytochemical investigation for the presence of various phytochemical constituents [11].

### Acute oral toxicity studies (LD<sub>50</sub>)

The acute toxicity of alcoholic extract of *P. spicigera* seeds were determined in wister rat fasted for 3 hour. (which examined that /DEL). The highest oral dose administered was 3g / kg body weight (which was equivalent to powder crude drug 18.95g / Kg of body weight). Up to 3g/Kg dose

levels no signs of toxicity appeared. The  $LD_{50}$  of the test extracts were calculated using AOT 425 software [10].

Oral toxicity: Not considered as toxic ( $DL_{50}$  oral/Rats > 30g/Kg body weight).

#### Preparation of drugs and chemicals

Ibuprofen (90mg  $Kg^{-1}$ ) Sigma, USA and acetylsalicylic acid (ASA) (10mg  $Kg^{-1}$ ) Sigma, USA were used as reference drugs and prepared by dissolving them in distilled water ( $dH_2O$ ). Other chemicals used were: Absolute alcohol, distilled water ( $dH_2O$ ), acetylsalicylic acid, Ibuprofen.

#### Experimental animals:

Wister rats (150-180g) were used to study anti-inflammatory activity. All these animals were maintained under standard husbandry protocol and conditions (light/dark period of 12 h light /dark and temperature  $25^{\circ}C \pm 3^{\circ}C$ ) with free access to food and water ad libitum and all experiments were carried out between 10 am to 5 pm daily [17].

#### Anti-inflammatory activity

Starved rats (48 h), weighing 150-180g having free access to drinking water were placed in separate single-single cages with raised bottom in order to avoid cannibalism and coprophagy [16]. The rats were randomly allotted to five groups containing five animals each [17] for the anti-inflammatory activity study and received distilled water ( $dH_2O$ ), 90mg  $kg^{-1}$  Ibuprofen or EPSS (33, 100 and 300 mg  $Kg^{-1}$ ), respectively, 1h prior to subjection to the test. All of the test solutions were administered in the volume of 10 ml  $Kg^{-1}$  body weight.

Table- 1 : Experimental groups and treatment given

Groups (n=5)	Treatment (Dose/Kg, p. o.)
Group 1	33 mg $Kg^{-1}$ EPSS
Group 2	100 mg $Kg^{-1}$ EPSS
Group 3	300 mg $Kg^{-1}$ EPSS
Group 4	90 mg $Kg^{-1}$ Ibuprofen
Group 5	Saline

According to Winter et. al.(1962) each five groups containing five rats received normal Saline, Ibuprofen, EPSS ( 33,100 and 300 mg  $Kg^{-1}$ ) followed 1 h later by the administration of 0.1 ml of 1% carrageenan suspension into

the rats right hind paw. Paw volume was measured before ( $V_0$ ) and 1,2,3,4 and 5 h ( $V_t$ ) following the carrageenan injection using a plethysmometer ( Model 7140, Ugo Basile, Italy ). The inflammation was measured by volume displaced by the paw between the final volume ( $V_t$ ) and the initial volume ( $V_0$ ). Mean of inflammation score for each animal was expressed by formula given:

$$\text{Percentage of anti-inflammation} = \frac{(V_t - V_0)_{\text{controlled}} - (V_t - V_0)_{\text{treated}}}{(V_t - V_0)_{\text{controlled}}} \times 100$$

#### Statistical Analysis

The results are expressed as mean  $\pm$  SEM. Statistical difference between means were determined by one way ANOVA followed by Dennett's post hoc test was used to analyze and compared data with  $P > 0.05$  as the limit of significance. (SEM = Standard Error of Mean).

### 3. RESULTS AND DISCUSSION

#### Preliminary phytochemical investigation:

he preliminary phytochemical screening with seeds extract of *Prosopis spicigera* revealed the presence of carbohydrates, amino acids and flavonoids in aqueous extract, alkaloids, flavonoids and tri-terpenoids in alcoholic extract and only steroids in petroleum ether extract [15,18].

#### Acute toxicity Study

Aqueous and alcoholic extracts up to a dose of 5000 mg/kg body wt and petroleum ether Extracts up to a dose of 2500 mg/kg were found to be safe[7].

#### The anti-inflammatory profile of the ethanolic *P. spicigera* supernatant ( EPSS )

All doses the EPSS significantly (  $p < 0.05$  ) inhibited the development of paw edema resulted from the carrageenan administration as shown in table 2 . The activity also occurred in a dose - dependent manner and started 1 h or their administration and lasted until the end of the experiment. The 10 mg  $kg^{-1}$ . Acetyl salicylic acid ( ASA ) also shows the pattern of activity as shown by \$ in table 2.

Table 2 :  
Effect of *P.spicigera* extract ( EPSS ) on Anti-inflammatory activity by the carrageenan-induced paw edema test in rats.

Treatment Groups (n=5)→ Mean increase In paw edema ±SEM (ml)/Time Interval (h) ↓	33mg Kg <sup>-1</sup> EPSS	100 mg Kg <sup>-1</sup> EPS	300 mg Kg <sup>-1</sup> buprofen	Saline	90mg Kg <sup>-1</sup>
1 h	0.40±0.025*	0.14±0.025*\$	0.12±0.02*	0.56±0.05	0.12±0.05*
2 h	0.38±0.025*\$	0.19±0.025*\$	0.16±0.02*	0.66±0.05	0.18±0.05*
3 h	0.36±0.03*\$	0.23±0.025*\$	0.18±0.02*	0.79±0.05	0.15±0.05*
4 h	0.31±0.03*\$	0.25±0.025*\$	0.16±0.02*\$	0.85±0.05	0.13±0.05*
5 h	0.29±0.03*\$	0.21±0.025*\$	0.15±0.02*	0.83±0.05	0.10±0.05*

The volume of hind paw edema was expressed as mean ± SEM,

\*Data differs significantly (  $p \leq 0.05$  ) when compared against the normal Saline-treated group.

\$Data differs significantly (  $p \leq 0.05$  ) when compared against ASA-treated group.

The present study indicated the ability of EPSS to exert anti-inflammatory activity in various animal models. The ability to inhibit chemically and thermally indicates as strong analgesics [5]. The abdominal constriction induced by the acetic acid were due to the release of cyclo- Oxyge-nase-synthesized prostacycline [1]. Which in turn lead to inflammatory pain with in the peritoneal cavity. [7]

The results indicated that *Prosopis spicigera* extracts produced anti-inflammatory effects Possessing antisecretory ,cytoprotective and proton pump inhibition mechanism[5,11].

The present study represented the potential of *Prosopis spicigera* seeds to exert anti-inflammatory activity especially the alcoholic extract.

#### 4. SUMMARY

In conclusion, the present study indicated the potential of *Prosopis spicigera* seeds extract has a potential anti-inflammatory activity especially the alcoholic extract and thus justify the folklore uses of the plant in treating pain and inflammation related ailments.

#### ACKNOWLEDGEMENT

The authors would like to thanks Dr. Ashok Kumar Srivastava Principal & Dr. Nagendra Swaroop, Honorable Secretary, Board of Management, D.B.S. (PG) College, Kanpur-06 (U.P.) and C.D.R.I. Lucknow (U.P.) & D.R.D.E. Gwalior (M.P.) for providing the facilities for the work and for their encouragement.

#### 5. REFERENCES

- [1]. Ballou, L.R., Botting, S., Zhang, J. and Vane, J.R.,2000. Nociception cyclooxygenase isozyme-deficient rats.Proc. Natl. Acad. Sci. USA, 97 (18), 10272-276.
- [2]. Begum N.A., Chaudhary D.N., Banerji J. and Das B.P., 2005. "J. Ind. Chem. Soc." Vol.82, pp.165-171, Feb.
- [3]. Bhardwaj et.al.,1980."Phytochemistry" jan.1979 ,18 (2) p.355-356, and19 (6) p.1269-70 .
- [4]. Chopra, R.N., Nayar, S.L. and Chopra, I.C., 1956. " Glossary of Indian Medical plants" Council of Scientific & Industrial Research, India. New Delhi, India. pp. 228-229.
- [5]. Gamache, D.A., Povlishook J.T. and Ellis, E.F., 1986, "Carrageenan-induced brain inflammation. Characterization of the model". J. Neurosurg., 65:679-685.
- [6]. Geissman, T.A. and Venkatraman, K.,1962. "chemistry of Flavonoid compounds". Ed. p.72 pergman press Oxford.
- [7]. Hunskaar S and Hole. K.,1987. The formalin test in mice:Dissocation between inflammatory and non-inflammatory pain. Pain, 30:103-114.
- [8]. Inuma M., Mosayoshi O.,Toshiyuki T. and Mizuo M.,1991. "J. of Natural Products" 54, (4) p. 1144.
- [9]. Jewer, K.et. al.,1976. "Phytochemistry" 15 (1) p. 238-240 jan.
- [10]. Kulkarni S.K.,1999. "Handbook of experimental Pharmacology."3rd. Ed. New Delhi. Vallabh Prakashan, p. 148-50.
- [11]. Khandelwal K.R.,2000. "Practical pharmacognosy techniques and experiment" 2nd edition Pune: Nirali prakashan 149-156.

- [12] Madan, B.R. et. al.,1972. "Ind. J. of Physiology and Pharmacology" Vol. 16 (2) pp.145-150, April.
- [13] Prakash P., Gupta N.,2005. "Ind. J. Physiol. Pharmacol." 49(2) 125-131.
- [14] Ray Sahelian,2003. "Indian J.Exp.Biol. "Nov 41(II) 1329-33 India.
- [15] Vogel, H.G and Vogel W.H.,1997. "Drug discovery and evaluation": Pharmacological assays. Lewisville, J.A. Majors Company, pp: 360-418.
- [16] Vogel H.G.,2002. "Drug Discovery and evaluation" II Ed. New York, Springer-Verlag, Berlin, Heidelberg. p. 867.
- [17] Winter, C.A., Risley, E. A.and Nuss, G.W.,1962."Carrageenin-induced edema in hind paws of the rats as a assays for anti-inflammatory drugs". Proc. Soc. Exp. Biol. Med., 111: 544-547.
- [18] Yadava, R.N.,2005. Novel biologically active flavonoidal constituent from *Smithia conferta* Sm. In: Proceedings of the 4th World Congress on Allelopathy. Eds. J. Harper, M., An, H. Wu, and J.H. Kent, International Allelopathy Society. The Regional Institute Ltd. Gosford NSW 2250, Charles Sturt University, Wagga Wagga, NSW, Australia. ISBN: 1864671688, August 21-26(2005).
- [19] Zimmerman, M.,1983. Ethical guidelines for investigations of experimental pain in conscious animals. Pain, 16: 109-110.